

Modifying the Order of Medication Search Results in an Electronic Health Record to Increase Physician Generic Prescribing Behavior

John Schrom, MPH, Paul Cohen, Sophie Krisch, Tamer Fakhouri, MD

Introduction

Prescription drug costs accounted for 10% of national health expenditure in the United States in 2016[1]. Generic drugs have the potential to eliminate significant spending when appropriately substituted, and are associated with greater patient adherence.

Malhotra et al showed that reengineering an e-prescribing interface to convert provider's selections of branded drugs to generic drugs significantly increased the proportion of generic prescriptions. However, this approach may not be appropriate for other workflows, such as when providers must choose between non-equivalent options.

We present a simple intervention whereby generic options are promoted within search results while still allowing providers to choose brand or generic options. The intervention significantly increased the rate of generic prescribing without requiring either changes to the design of the ordering module or dedicated provider communication.

Methods

Intervention

The home-grown electronic health record for a national primary care clinic system allows for search results to be reordered based on matching against arbitrarily defined medication synonyms. Branded medication names were listed as synonyms for their respective generic medications, prompting the generic to be promoted when a physician searches for its corresponding branded medication.

This change was made for four medications:

- valacyclovir (Valtrex),
- escitalopram oxalate (Lexapro),
- amoxicillin-potassium clavulanate (Augmentin),
- atorvastatin (Lipitor).

Evaluation

New prescriptions were tracked for two months before and after the launch of the intervention. Data was analyzed using Pearson's chi-squared test, with the SciPy open source package for Python, version 0.19.1.

MEDICATIONS

Medication Name


aug

- amoxicillin-pot clavulanate (Augmentin)
- Augmentin amoxicillin-pot clavulanate
- Augmentin ES-600 amoxicillin-pot clavulanate
- Augmentin XR amoxicillin-pot clavulanate
- betamethasone, augmented topical

Results

In the study period, 58,384 prescriptions were made for the branded or generic versions of the test medications. Each medication showed a significant increase in the proportion of generic prescriptions ($p < 0.0001$). There was no change in prescriptions with the "dispense as written" indicator.

	Before	After
valacyclovir	33%	61%
escitalopram oxalate	29%	55%
amoxicillin-potassium clavulanate	5%	65%
atorvastatin	75%	87%



Conclusions

Changing the position of equivalent generic medications in search results significantly increases the generic prescribing rate without subverting physician intent. This suggests that minor user experience changes, such as the list ordering, can be used to nudge clinicians into making higher quality and better value decisions with their patients.

References

1. CMS. National Health Expenditures 2016 Highlights. Retrieved from: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>
2. Generic Pharmaceutical Association. Generic drug savings in the US. 2015. www.gphaonline.org/media/wysiwyg/PDF/GPhA_Savings_Report_2015.pdf
3. Shrank WH, Hoang T, Ettner SL, et al. The implications of choice: prescribing generic or preferred pharmaceuticals improves medication adherence for chronic conditions. Arch Intern Med 2006;166:332-7. 10.1001/archinte.166.3.332
4. Malhotra S, Cheriff AD, Gossey JT, Cole CL, Kaushal R, Ancker JS. Effects of an e-prescribing interface redesign on rates of generic drug prescribing: exploiting default options. J Am Med Inform Assoc. 2016 Sep;23(5):891-8